

**NEUTRON**  
CLEAN SMARTER. SMELL BETTER.

READY TO USE • LAVENDER SCENT

# TRIPLE Threat Defense

**DISINFECTANT • BACTERICIDAL  
 †VIRUCIDE • TUBERCULOCIDAL  
 MILDREWSTAT (on hard inanimate surfaces)  
 FUNGICIDE (against pathogenic fungi)  
 CLEANER • DEODORIZER**

Kills SARS-CoV-2 (COVID 19 virus) on hard non-porous surfaces

Meets OSHA Bloodborne Pathogen Standard for HIV, HBV &amp; HCV

ACTIVE INGREDIENTS

n-Alkyl (60% C <sub>14</sub> , 30% C <sub>16</sub> , 5% C <sub>12</sub> , 5% C <sub>18</sub> )	
dimethyl benzyl ammonium chlorides.....	0.105%
n-Alkyl (68% C <sub>12</sub> , 32% C <sub>14</sub> )	
dimethyl ethylbenzyl ammonium chlorides.....	0.105%
<b>INERT INGREDIENTS.....</b>	<b>99.790%</b>
<b>TOTAL.....</b>	<b>100.000%</b>

EPA REG. NO. 1839-83-42666

EPA EST. NO. 70799-0H-2

## KEEP OUT OF REACH OF CHILDREN CAUTION

See side panels for additional precautionary statements.

### FIRST AID

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. **If in eyes:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice. **If swallowed:** Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person. **If inhaled:** Move person to fresh air. If person is not breathing, call 911 or an ambulance then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice. **NOTE TO PHYSICIAN:** Probable mucosal damage may contraindicate the use of gastric lavage.

NET CONTENTS: 1 Quart (32 FL Oz.) (0.946 liters)

This product is designed specifically as a general non-acid cleaner and disinfectant for use on hard, non-porous surfaces in hospitals, nursing homes, schools, food processing plants, food service establishments, transportation terminals, office buildings, manufacturing facilities, lodging establishments, retail businesses, veterinary clinics, pet shops, animal life science laboratories and athlete/recreational facilities where housekeeping is of prime importance in controlling the hazard of cross contamination. It is formulated to disinfect hard, non-porous, inanimate environmental surfaces: floors, walls, metal surfaces, stainless steel surfaces, glazed porcelain, glazed ceramic tile, plastic surfaces, chrome, brass, copper, laminated surfaces, baked enamel surfaces, bathrooms, shower stalls, bathtubs, and cabinets. **For plastic and painted surfaces, spot test on an inconspicuous area before use.** May be used in the kitchen on counters, appliances and stovetops. A rinse with potable water is required for surfaces in direct contact with food. This product has demonstrated effectiveness against Influenza A Virus and is expected to inactivate all Influenza A Viruses including Pandemic 2009 H1N1 Influenza A Virus. Triple Threat Defense kills 99.9%\* of SARS-CoV-2, which causes COVID-19\*\* on hard non-porous surfaces in 1 minute.

**EFFICACY TESTS HAVE DEMONSTRATED THAT THIS PRODUCT IS AN EFFECTIVE BACTERICIDE, FUNGICIDE, AND VIRUCIDE† IN THE PRESENCE OF ORGANIC SOIL (5% BLOOD SERUM).**

### DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

This product is not to be used as a terminal sterilant/high-level disinfectant on any surface or instrument that (1) is introduced directly into the human body, either into or in contact with the bloodstream or normally sterile areas of the body, or (2) contacts intact mucous membranes but which does not ordinarily penetrate the blood barrier or otherwise enter normally sterile areas of the body. **DISINFECTION, DEODORIZING AND CLEANING -** Remove visible soil prior to application of the product. Hold container six to eight inches from surface to be treated. Spray area until it is covered with the solution. Allow product to penetrate and the surface to remain wet for **3 minutes** (for Bloodborne Pathogens: allow surface to remain wet for HIV-1 **1 minute** and for HBV and HCV **1 minute**). No scrubbing is necessary. Wipe off with a clean cloth, mop or sponge. The product will not leave grit or soap scum.

**One-step\* cleaning and disinfection of toilet bowls and urinals:** Remove water from bowl by forcing water over trap with a swab mop applicator. Press swab mop against side of the bowl to remove excess water from applicator. Apply 1 to 2 ounces of this product evenly, allowing sufficient time for product to cover entire surface of bowl for maximum cleaning. Swab bowl completely using a scrub brush or toilet mop, making sure to get under the rim. Let stand for **3 minutes** (for Bloodborne Pathogens: allow surface to remain wet for HIV-1 **1 minute** and for HBV and HCV **1 minute**) and flush. If water cannot be removed from toilet, use product on the external surface and above waterline only.

**To disinfect food processing premises:** Before using this product, food products and packaging materials must be removed or carefully protected. For floors, walls and storage areas, hold container six to eight inches from the surface to be treated and spray area until it is covered with the solution. Allow product to penetrate and remain wet for **3 minutes** (for Bloodborne Pathogens: allow surface to remain wet for HIV-1 **1 minute** and for HBV and HCV **1 minute**). Wipe off with a clean cloth, mop or sponge and thoroughly rinse all surfaces in the area with potable water. Visibly soiled areas must be pre-cleaned before using this product.

**To disinfect food service establishment food contact surfaces:** Before using this product, food products and packaging materials must be removed or carefully protected. For countertops, appliances, tables, hold container six to eight inches from the surface to be treated and spray area until it is covered with the solution. Allow product to penetrate and remain wet for **3 minutes** (for Bloodborne Pathogens: allow surface to remain wet for HIV-1 **1 minute** and for HBV and HCV **1 minute**). Wipe off with a clean cloth, mop or sponge and thoroughly rinse all surfaces in the area with potable water. Visibly soiled areas must be pre-cleaned before using this product. Do not use on eating utensils, glassware and cookware.

**Disinfection of animal quarters and kennels:** Before using this product, remove all animals and feed from premises. Remove all litter and droppings from floors, walls and surfaces of facilities occupied or traversed by animals. Empty all troughs, racks, and other feeding and watering appliances. Thoroughly clean all surfaces with soap or detergent and rinse with water. Apply this product for disinfection of pre-cleaned animal quarters and kennels. Saturate the surfaces with this product for a period of **3 minutes** (for Bovine Viral Diarrhea Virus (BVDV) **1 minute**) (for Feline Calicivirus and Rabies virus **30 seconds**) (for Canine Parvovirus **10 minutes**). Ventilate building and other closed spaces. Do not house animals or employ equipment until treatment has been absorbed, set or dried. All tested equipment that will contact feed or drinking water must be rinsed with potable water before reuse.

Using AOAC test methods under Good Laboratory Practices, [GLPs], in the presence of 5% blood serum and a **3 minute** contact time, unless otherwise noted, this product kills the following organisms on hard non-porous inanimate surfaces:

**BACTERICIDAL ACTIVITY:** Pseudomonas aeruginosa, Staphylococcus aureus, Salmonella (choleraesuis) enterica, Escherichia coli, Escherichia coli O157:H7, Streptococcus pyogenes (Necrotizing Fasciitis-Group A), Listeria monocytogenes, Yersinia enterocolitica, Enterococcus faecium, Corynebacterium ammoniagenes, Salmonella (typhi) enterica.

Claim Qualifier: \*when used according to disinfection directions

Claim Qualifier: \*\*SARS-Related Coronavirus 2

**ANTIBIOTIC-RESISTANT BACTERICIDAL ACTIVITY:** Methicillin resistant Staphylococcus aureus (MRSA), Vancomycin resistant Enterococcus faecalis (VRE) (ATCC 51575), Vancomycin intermediate resistant Staphylococcus aureus (VISA) (CDC Isolate 99287), Methicillin resistant Staphylococcus epidermidis (MRSE), Community Associated Methicillin resistant Staphylococcus aureus (CA-MRSA) (NRS 123) Genotype USA400, Community Associated Methicillin resistant Staphylococcus aureus (CA-MRSA) (NRS 384) Genotype USA300.

**TUBERCULOCIDAL ACTIVITY -** This product exhibits disinfectant efficacy against Mycobacterium tuberculosis BCG (Mycobacterium bovis) at 20 degrees Centigrade when the treated surface is allowed to remain wet for **5 minutes**.

**VIRUCIDAL ACTIVITY -** This product kills on hard, non-porous inanimate surfaces when allowed to remain wet for a **10 minute** contact time against: Canine Parvovirus, Hepatitis A Virus (HAV) (University of Ottawa), Poliovirus Type 1. **This product, when used on environmental, inanimate hard surfaces exhibits virucidal activity against:** Human Coronavirus, SARS associated Coronavirus, Avian Influenza A strain H3N2 and Avian Influenza A strain H9N2 for **2 minutes**; Paramyxovirus (Mumps), Rhinovirus type 39 and Rotavirus (University of Ottawa) for **3 minutes**; HIV-1 (associated with AIDS) (Advanced Biotechnologies, Inc.), SARS-Related Coronavirus 2 (SARS-CoV-2) for **1 minute**; Hepatitis B Virus (HBV) (Hepadnavirus Testing, Inc.), Hepatitis C Virus (HCV) and Bovine Viral Diarrhea Virus (BVDV) (NVSL) when the treated surface is allowed to remain wet for **1 minute**; and Norovirus, Feline Calicivirus and Rabies Virus for **30 seconds**.

**KILLS HIV-1, HBV AND HCV ON PRE-CLEANED ENVIRONMENTAL SURFACES/OBJECTS PREVIOUSLY SOILED WITH BLOOD/BODY FLUIDS** in health care settings (Hospitals, Nursing Homes) or other settings in which there is an expected likelihood of soiling of inanimate surfaces/objects with blood or body fluids, and in which the surfaces/objects likely to be soiled with blood or body fluids can be associated with the potential for transmission of Human Immunodeficiency Virus Type 1 (HIV-1) (associated with AIDS), Hepatitis B Virus (HBV), and Hepatitis C Virus (HCV).

**SPECIAL INSTRUCTIONS FOR CLEANING AND DECONTAMINATION AGAINST HIV-1, HBV AND HCV ON SURFACES/OBJECTS SOILED WITH BLOOD/BODY FLUIDS. PERSONAL PROTECTION:** When handling items soiled with blood or body fluids use disposable latex gloves, gowns, masks, and eye coverings. **CLEANING PROCEDURES:** Blood and other body fluids must be thoroughly cleaned from surfaces and objects before application of this product. **CONTACT TIME:** Allow surface to remain wet for HBV and HCV **1 minute**, for HIV-1 **1 minute**. **These contact times will not control other common types of viruses and bacteria. DISPOSAL OF INFECTIOUS MATERIALS:** Blood and other body fluids must be autoclaved and disposed of according to local regulations for infectious waste disposal.

**FUNGICIDAL ACTIVITY -** This product is fungicidal against the pathogenic fungus, Trichophyton interdigitale, formerly Trichophyton mentagrophytes (Athlete's Foot Fungus), when used with a 10 minute contact time on clean, non-porous, hard surfaces as found in bathrooms, shower stalls, locker rooms, or other areas commonly contacted by bare feet. **MILDREWSTAT -** To control mold and mildew on pre-cleaned, hard, non-porous surfaces spray surface to be treated making sure to wet completely. Let air dry. Repeat application at weekly intervals or when mildew growth appears.

## PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

**KEEP OUT OF REACH OF CHILDREN.** Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum or using tobacco.

## STORAGE AND DISPOSAL

**DO NOT CONTAMINATE WATER, FOOD, OR FEED BY STORAGE OR DISPOSAL.**

**Pesticide Storage -** Store in a dry place no lower in temperature than 50° F or higher than 120° F. **Pesticide Disposal -** Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance. **Container Disposal -** Nonrefillable container. Do not reuse or refill container. Clean container promptly after emptying. Triple rinse as follows: Fill container ¼ full with water and recap. Agitate vigorously. Follow Pesticide Disposal instructions for rinsate disposal. Drain for 10 seconds after the flow begins to drip. Repeat procedure two more times. Offer for recycling or reconditioning, if available. If not available, puncture and dispose in a sanitary landfill.