

SPORICIDAL PERFORMANCE AGAINST C. DIFFICILE SPORES: This product kills and/or inactivates the following spore in 2 minutes at 4 oz. per gal. of 400 ppm hard water (1844 ppm active PAA) on hard, non-porous surfaces: Clostridium difficile.

TUBERCULOCIDAL PERFORMANCE: This product kills the following mycobacteria in 10 minutes at 4 oz. per gal. of 400 ppm hard water (1844 ppm active PAA) and 5% soil at 21°C, on hard, non-porous surfaces: Mycobacterium bovis (Tb).

DISINFECTANT PERFORMANCE: This product kills the following bacteria in 2 minutes at 2 oz. per gal. of 400 ppm hard water (922 ppm active PAA) and 5% soil, on hard, non-porous surfaces: Acinetobacter baumannii, Bordetella pertussis, Enterococcus faecalis Vancomycin Resistant (VRE), Escherichia coli, Escherichia coli (Extended Spectrum B-Lactamase) (ESBL), Klebsiella pneumoniae, Klebsiella pneumoniae Carbapenem Resistant, Proteus mirabilis, Pseudomonas aeruginosa, Salmonella enterica, Staphylococcus aureus, Staphylococcus aureus Community Acquired Methicillin Resistant (CA-MRSA), Staphylococcus aureus Methicillin Resistant (MRSA), Staphylococcus aureus Vancomycin Intermediate Resistant (VISA), Streptococcus pneumoniae, Streptococcus pyogenes.

This product kills the following bacteria in 10 minutes at 2 oz. per 5 gal. of 400 ppm hard water (184 ppm active PAA) and 5% soil, on hard, non-porous surfaces: Bordetella bronchiseptica, Campylobacter jejuni, Corynebacterium ammonigenes, Enterococcus faecalis Vancomycin Resistant (VRE), Escherichia coli O157:H7, Klebsiella pneumoniae, Listeria monocytogenes, Pseudomonas aeruginosa, Salmonella enterica, Salmonella typhi, Shigella sonnei, Staphylococcus aureus, Staphylococcus aureus Community Acquired Methicillin Resistant (CA MRSA), Staphylococcus aureus Vancomycin Intermediate Resistant (VISA).

VIROCIDAL PERFORMANCE: This product kills the following viruses in 2 minutes at 4 oz. per gal. of 400 ppm hard water (1844 ppm active PAA) and 5% soil, on hard, non-porous surfaces: Adenovirus Type 5, Hepatitis B Virus † (HBV), Hepatitis C Virus † (HCV), Herpes Simplex Type 1 Virus, Herpes Simplex Type 2 Virus, Human Immunodeficiency Virus Type 1 (HIV-1), Influenza A Virus, Norovirus (Nonwalk-like Virus), Respiratory Syncytial Virus, Rhinovirus Type 37, Rotavirus, Vaccinia Virus. †Indicates a 5-minute contact time is required for this claim.

FOOD CONTACT SANITIZING PERFORMANCE: This product is an effective food contact surface sanitizer in 1 minute at 2 oz. per 6 gal. of 500 ppm hard water (154 ppm active PAA) on hard, non-porous surfaces: Aeromonas hydrophila, Clostridium perfringens - vegetative, Enterobacter sakazakii, Escherichia coli, Klebsiella pneumoniae, Listeria monocytogenes, Salmonella enterica, Salmonella typhi, Shigella dysenteriae, Shigella sonnei, Staphylococcus aureus, Yersinia enterocolitica.

FUNGICIDAL PERFORMANCE: This product is effective against the following organisms in 2 minutes at 4 oz. per gal. of 400 ppm hard water (1844 ppm active PAA) and 5% soil, on hard, non-porous surfaces: Candida albicans, Trichophyton interdigitale (Athlete's foot fungus).

ANIMAL PREMISE VIROCIDAL PERFORMANCE: This product kills the following viruses in 2 minutes at 4 oz. per gal. of 400 ppm hard water (1844 ppm active PAA) and 5% soil, on hard, non-porous surfaces: Canine Parvovirus †, Murine Norovirus, Porcine Epidemic Diarrhea Virus. †Indicates a 5-minute contact time is required for this claim.

This product kills the following viruses in 10 minutes at 2 oz. per 5 gal. of 400 ppm hard water (184 ppm active PAA) and 5% soil, on hard, non-porous surfaces: Avian Adenovirus, Avian Infectious Bronchitis Virus, Avian Influenza A Virus, Infectious Bursal Disease Virus, Infectious Laryngotracheitis Virus, Newcastle Disease Virus, Porcine Rotavirus, Pseudorabies Virus, Transmissible Gastroenteritis Virus, Vesicular Stomatitis Virus.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Before using this product in federally inspected meat and poultry food processing plants and dairies, food products and packaging materials must be removed from the room or carefully protected.

This product is not for use on medical device surfaces.

AGRICULTURAL USE REQUIREMENTS: Use this product only in accordance with its labeling and with the Worker Protection Standard, 40CFR170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), restricted-entry interval, and notification to workers. The requirements in this section (box) only apply to uses of this product that are covered by the Worker Protection Standard. Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 48 hours.

Notify workers of the application by warning them orally and by posting warning signs at entrances to treated areas. PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is: coveralls worn over long sleeved shirt and long pants, waterproof gloves, chemical-resistant shoes plus socks, and protective eyewear.

SOLD BY:

State Industrial Products
5915 Landerbrook Drive
Mayfield Heights, OH 44124
To Order Call: 1-866-747-2229
www.statechemical.com

Puerto Rico
Royal Industrial Park, Bldg "M"
Local #53, Carr 869, km 1.5 Palmes
Cataño, PR. 00962
To Order Call: 787-275-3185

J-17

81N11 125751BL
©2018-2021 State Industrial Products

FOR USE AS A SPORICIDE AGAINST Clostridium difficile SPORES: 1. Pre-clean surfaces. Fecal matter/waste must be thoroughly cleaned from surfaces/objects before disinfection by application with a clean cloth, mop, and/or sponge saturated with the disinfectant product. This cleaning may be accomplished with any cleaning solution, including this product. Cleaning is to include vigorous wiping and/or scrubbing, until all visible soil is removed. Special attention is needed for high-touch surfaces. Surfaces in patient rooms are to be cleaned in an appropriate manner, such as from right to left or left to right, on horizontal surfaces, and top to bottom, on vertical surfaces, to minimize the spreading of the spores. Restrooms are to be cleaned last. Do not reuse soiled cloths. 2. Apply use solution of 4 oz. of this product per gal. of water to disinfect hard, non-porous surfaces with a sponge, brush, cloth, mop, by immersion, mechanical spray device, hand pump trigger spray device. For spray applications, spray 6-8 inches from surface. Do not breathe spray. 3. Treated surfaces must remain wet for 2 minutes. 4. Wipe dry with a clean cloth or allow to air dry. 5. Prepare a fresh solution daily or when visibly dirty. Note: Materials used in the cleaning process that may contain feces/wastes are to be disposed of immediately in accordance with local regulations for infectious materials disposal.

FOR USE AS A TUBERCULOCIDE: 1. Pre-clean heavily soiled areas. 2. Apply use solution of 4 oz. of this product per gal. of water (1844 ppm active PAA) at 21°C to disinfect hard, non-porous surfaces with a sponge, brush, cloth, mop, by immersion, auto scrubber, mechanical spray device, hand pump trigger spray device. For spray applications, spray 6-8 inches from surface. Do not breathe spray. 3. Treated surfaces must remain wet for 10 minutes. 4. Wipe dry with a clean cloth or allow to air dry. 5. Prepare a fresh solution daily or when visibly dirty.

FOR USE AS A VIROCIDE†, FUNGICIDE: 1. Pre-clean heavily soiled areas. 2. Apply use solution of 4 oz. of this product per gal. of water (1844 ppm active PAA) to disinfect hard, non-porous surfaces with a sponge, brush, cloth, mop, by immersion, auto scrubber, mechanical spray device, hand pump trigger spray device. For spray applications, spray 6-8 inches from surface. Do not breathe spray. 3. Treated surfaces, must remain wet for 2 minutes. 4. Wipe dry with a clean cloth or allow to air dry. 5. Prepare a fresh solution daily or when visibly dirty.

FOR USE AS A HOSPITAL DISINFECTANT: 1. Pre-clean heavily soiled areas. 2. Apply use solution of 2 oz. per 5 gal. of water (184 ppm active PAA) to disinfect hard, non-porous surfaces with a sponge, brush, cloth, mop, by immersion, auto scrubber, mechanical spray device, hand pump trigger spray device. For spray applications, spray 6-8 inches from surface. Do not breathe spray. 3. Treated surfaces, must remain wet for 10 minutes. 4. Wipe dry with a clean cloth or allow to air dry. 5. Prepare a fresh solution daily or when visibly dirty.

FOR USE AS A ONE-STEP, GENERAL, HOSPITAL DISINFECTANT: 1. Pre-clean heavily soiled areas. 2. Apply use solution of 2 oz. of this product per gal. of water (400 ppm active PAA) to disinfect hard, non-porous surfaces with a sponge, brush, cloth, mop, by immersion, auto scrubber, mechanical spray device, hand pump trigger spray device. For spray applications, spray 6-8 inches from surface. Do not breathe spray. 3. Treated surfaces, must remain wet for 2 minutes. 4. Wipe dry with a clean cloth or allow to air dry. 5. Prepare a fresh solution daily or when visibly dirty.

***KILLS HIV-1, HBV AND HCV ON PRE-CLEANED HARD, NON-POROUS SURFACES/OBJECTS PREVIOUSLY SOILED WITH BLOOD/BODY FLUIDS** in health care settings or other settings in which there is an expected likelihood of soiling of hard, non-porous surfaces/objects with blood or body fluids and in which the surfaces/objects likely to be soiled with blood or body fluids can be associated with the potential for transmission of Human Immunodeficiency Virus Type 1 (HIV-1), Hepatitis B Virus (HBV) and Hepatitis C Virus (HCV).

SPECIAL INSTRUCTIONS FOR CLEANING AND DECONTAMINATION AGAINST HIV-1, HBV AND HCV ON SURFACES/OBJECTS SOILED WITH BLOOD/BODY FLUIDS.

Personal Protection: Wear protective latex gloves, gowns, masks and eye protection. **Cleaning Procedure:** Blood and other body fluids must be thoroughly cleaned from hard, non-porous surfaces and objects before application of this product. **Disposal of Infectious Materials:** Blood and other body fluids, cleaning materials and clothing must be autoclaved and disposed of according to federal, state and local regulations for infectious waste disposal. **Contact Time:** † Allow solution to remain wet for 2 minutes for HIV-1, 5 minutes for HBV and HCV at 4 oz. of this product per gal. of water. **FOOD CONTACT SURFACE SANITIZING DIRECTIONS:** Prior to application, remove gross food particles and soil by a pre-rinse or pre-spray and when necessary, pre-soak. Then thoroughly wash or flush objects with a good detergent or compatible cleaner, followed by a potable water rinse before applications of the sanitizing solution.

SURFACES IN FOOD PROCESSING LOCATIONS, DAIRIES, RESTAURANTS AND BARS: Immerse pre-cleaned glassware, dishes, silverware, cooking utensils and other similar size food processing equipment in a solution of 2 oz. of this product per 6 gal. of water (0.34 oz. of this product per gal. of water)(154 ppm active PAA) (or equivalent use dilution) for at least 1 minute. Allow sanitized surfaces to adequately drain and then air dry before contact with food. Do not rinse. For articles too large for immersing, apply a use solution of 2 oz. of this product per 6 gal. of water (0.34 oz. of this product per gal. of water)(154 ppm active PAA) (or equivalent use dilution) to sanitize hard, non-porous food contact surfaces with a brush, cloth, mop, sponge, auto scrubber, mechanical spray device, hand pump trigger spray device. For spray applications, spray 6-8 inches from surface. Do not breathe spray. Surfaces must remain wet for at least 1 minute. Allow sanitized surfaces to adequately drain and then air dry before contact with food. Do not rinse. Prepare a fresh solution daily or when visibly dirty. For mechanical application, use solution must not be reused for sanitizing applications, but may be used for other purposes such as cleaning.

CLEAN-IN-PLACE (CIP) METHOD FOR DAIRY, DAIRY FARM AND FOOD PROCESSING FACILITIES: 1. Thoroughly flush, clean and potable water rinse the system. 2. Prepare required volume of sanitizer solution needed by diluting 0.34 oz. of this product per gal. of water (154 ppm active PPA) (or equivalent use dilution). 3. To sanitize entire system by circulation methods, run pumps for at least 2 minutes to thoroughly wet and sanitize all parts of the system.

State
Since 1911

STATE
POWER₂
DC

- Effective one-step disinfectant-cleaner for use in hospitals, long term care facilities, and other healthcare settings.
- Is a food contact surface sanitizer on hard, non-porous surfaces.
- Effective in 2 minutes against C. difficile spores.

ACTIVE INGREDIENTS:

Hydrogen Peroxide.....	27.3%
Peroxyacetic Acid.....	5.9%
OTHER INGREDIENTS:.....	66.8%
TOTAL:	100.0%

EPA Reg. No. 10324-214-70799
EPA Est. No. 35772-IL-01

KEEP OUT OF REACH OF CHILDREN DANGER PELIGRO

See side panel for additional precautionary statements.

FIRST AID: In case of emergency, call a poison control center or doctor for treatment advice. Have the product container or label with you when calling a poison control center or doctor, or going for treatment. **IF IN EYES:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. **IF ON SKIN:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. **IF SWALLOWED:** Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person. Call a poison control center or doctor immediately for treatment advice. **IF INHALED:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible.

NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage.

NET CONTENTS/CONTENU NET: 1 U.S. Gallon (3.78L)

www.statechemical.com

Product #125751

CLEANING AND DISINFECTING HARD, NON-POROUS SURFACES ON PERSONAL PROTECTIVE EQUIPMENT (RESPIRATORS): Pre-clean equipment, if heavily soiled to ensure proper surface contact. Add 2 oz. of this product per gal. of water (922 ppm active PAA). Gently mix for uniform use solution. Apply use solution to surfaces of the respirator with a sponge, brush, cloth, by immersion, mechanical spray device, hand pump trigger spray device, coarse trigger spray device. For spray applications, spray 6-8 inches from surface. Do not breathe spray. Rub with brush, cloth, or sponge. Treated surfaces must remain wet for 2 minutes. Remove excess solution from equipment prior to storage. The user must comply with all OSHA regulations for cleaning respiratory protection equipment (29 CFR § 1910.134). Prepare a fresh solution daily or when visibly dirty.

ANIMAL PREMISES: Prior to use of this product, remove all animals and feed from premises, animal transportation vehicles, and enclosures, coops, crates, kennels, stables. Remove all litter, droppings and manure from floors, walls and surfaces of barns, pens, stalls, chutes and other surfaces of facilities and fixtures occupied or traversed by animals. Empty all troughs, racks and other feeding and watering appliances. Thoroughly clean surfaces with soap or detergent and rinse with water.

FOR USE AS AN ANIMAL PREMISE DISINFECTANT/VIROCIDE†: For heavily soiled areas, a pre-cleaning step is required. Apply a use solution of 2 oz. of this product per 5 gal. of water (184 ppm active PAA) to disinfect hard, non-porous surfaces with a sponge, brush, cloth, by immersion, mechanical spray device, hand pump trigger spray device. For spray applications, spray 6-8 inches from surface. Do not breathe spray. Immerse all halters and other types of equipment used in handling and restraining animals, as well as forks, shovels, and scrapers used for removing litter and manure in the use solution. Treated surfaces must remain wet for 10 minutes. Ventilate buildings, coops and other closed spaces. Do not house animals or employ equipment until treatment has been absorbed, set or dried. Thoroughly scrub all treated feed racks, troughs, automatic feeders, fountains and waterers and other treated equipment which can contact food or water with soap or detergent, and rinse with potable water before reuse.

FOR USE AS AN ANIMAL PREMISE DISINFECTANT/VIROCIDE†: For heavily soiled areas, a pre-cleaning step is required. Apply a use solution of 4 oz. of this product per gal. of water (1844 ppm active PAA) to disinfect hard, non-porous surfaces with a sponge, brush, cloth, by immersion, mechanical spray device, hand pump trigger spray device. For spray applications, spray 6-8 inches from surface. Do not breathe spray. Immerse all halters and other types of equipment used in handling and restraining animals, as well as forks, shovels, and scrapers used for removing litter and manure in the use solution. Treated surfaces must remain wet for 2 minutes. Ventilate buildings, coops and other closed spaces. Do not house animals or employ equipment until treatment has been absorbed, set or dried. Thoroughly scrub all treated feed racks, troughs, automatic feeders, fountains and waterers and other treated equipment which can contact food or water with soap or detergent, and rinse with potable water before reuse.

† For use against Canine Parvovirus, prepare a use solution of 4 oz. of this product per gal. of water at 5 minutes. **AGRICULTURAL OR HORTICULTURAL USES:** This product must never be mixed or combined with any other pesticide or fertilizer. Upon soil contact this product decomposes rapidly to oxygen, carbon dioxide and water. The product is harmful to fish if exposed on a continuous basis at concentrations of 0.5 ppm or more of PAA. Meter this product into pressurized pipes using a plastic or stainless steel injection/backflow device installed for enough upstream from the target equipment to ensure thorough mixing. For open flowing bodies of water, apply this product as far upstream as possible to allow adequate mixing prior to the flow entering any larger body of water. If open pouring of this product is required pour product as close to the surface of the water as possible to reduce odor exposure. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protect ed handlers may be in the area during application. For any requirement specific to your State and Tribe, consult the State/Tribal agency responsible for pesticide regulation.

FOLIAR SPRAY TREATMENT IN GREENHOUSES (Not for use in CA.): This product works immediately on contact with any plant surface for control/suppression of fungi. Apply this product to ornamentals, bedding plants, flowering plants, shrubs, and trees. To ensure that this fungicide is effective, thorough coverage and wetting of the foliage is necessary. **Initial Application:** 1. Use 2/3 - 1 1/3 oz. of this product per gal. of clean water (307 - 614 ppm active PAA). Do not reuse already mixed solution. Make fresh solution at least daily or when use solution becomes visibly dirty, soiled, or diluted. 2. Spray, mist or fog plants in the early morning or late evening. Do not breathe spray. 3. Thoroughly wet all surfaces of plant including upper and lower foliage, stems, branches, and stalks to ensure full contact with plant and flower tissue. 4. Apply for one to three consecutive days and then follow directions for preventative treatment after the initial application.

Weekly Treatment: 1. Use 0.14 - 0.23 oz. of this product per gal. of clean water (64-106 ppm active PAA) (or equivalent use-dilution). 2. Spray, mist or fog plants. Do not breathe spray. 3. Thoroughly wet all surfaces of plant including upper and lower foliage, stems, branches and stalks to ensure full contact with plant and flower tissue. 4. Spray every five to seven days as a Preventive treatment. 5. At the first sign of disease, spray daily with a dilution of 2/3 - 1 1/3 oz. of this product per gal. of water (or equivalent use-dilution) for three consecutive days and then resume weekly Preventive Treatment.

NON-AGRICULTURAL USE REQUIREMENTS: The requirements in this section apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses. Keep children and pets out of the treated area until sprays have dried.

FRUIT AND VEGETABLE WATER TREATMENT: This product is used to help control spoilage or decay-causing bacteria and fungi in water or ice that contacts raw unprocessed fruits and vegetables. The commodity must be continuously sprayed using coarse spray, or submerged using a solution containing 1 oz. of this product per 20 gal. of water (23 ppm active PAA) (or equivalent use-dilution) for a minimum contact time of 30 seconds. Adjust dose as necessary to maintain no more than 80 ppm active PAA. Remove excess water or allow to drain. If using the submersion method, replace with a fresh solution at least daily, or when solution becomes visibly soiled. A final potable water rinse is not required.

TREATMENT OF FRUIT AND VEGETABLE PROCESSING WATERS: Use this product for the treatment of waters used in the processing of raw fruits and vegetables. Mix this product with water either batch-wise or continuously at a rate of 60-195 oz. of this product per 1,000 gal. of water (28-90 ppm active PAA) (or equivalent use-dilution). The fruits and vegetables can be sprayed or submerged in the resulting solution for a minimum contact time of 30 seconds, followed by adequate draining. At this use-dilution, this product will control the growth of spoilage and decay causing non-public health organisms in process waters and on the surface of fresh cut or post-harvest fruits and vegetables. This product is not allowed to be used for control of any public health organism on fruit and vegetable surfaces.

POST-HARVEST SPRAY TREATMENT: Use this product to prevent bacterial and fungal diseases on post-harvest fruits and vegetables. Mix 0.31 - 0.51 oz. of this product per gal. of clean water. Spray fruit or vegetables to the point of runoff using a mechanical spray device, hand pump trigger spray device. For spray applications, spray 6-8 inches from surface. Do not breathe spray.

FOAM CLEANING OF FOOD AND NON-FOOD CONTACT SURFACES: For cleaning procedures, this product may be added to Macat® AO-12 (amine oxidant) and foamed on hard, non-porous equipment surfaces using foam generating equipment. The resilient foam blend can be used on equipment, floors, walls, ceilings, drains, etc. and should be left on the surface for a minimum of 1 minute. To mix manually or mechanically blend 1 - 6.1 oz. of this product and 6 - 12 oz. of Macat® AO-12 (foam additive) per 6 gal. of water. The dilution water must not exceed 150° F. On food contact surfaces do not exceed 6.1 oz. of this product per 6 gal. of water.

FOR USE AS A GENERAL CLEANER AND/OR DEODORIZER: Apply a use solution of 2 oz. of this product per gal. of water to hard, non-porous surfaces. Wipe up excess liquid and allow to air dry. For heavy-duty use, add 4 oz. of this product per gal. of water to clean hard, non-porous surfaces.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

DANGER. CORROSIVE. Causes irreversible eye damage and skin burns. Harmful if swallowed. May be fatal if inhaled. Do not get into eyes, on skin or on clothing. Do not breathe vapors or spray mist. Wear goggles or face shield and chemical-resistant gloves and protective clothing when handling. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove contaminated clothing and wash clothing before reuse.

PERSONAL PROTECTIVE EQUIPMENT

Applicators and other handlers must wear coveralls worn over long-sleeved shirt and long pants, waterproof gloves, chemical-resistant footwear and socks, protective eyewear, chemical-resistant headgear when using this product for algae control in overhead watering system and chemical-resistant apron when mixing, loading or cleaning equipment. Discard clothing and other absorbent materials that have been drenched or heavily contaminated with the product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing. Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to birds, fish and aquatic invertebrates.

PHYSICAL OR CHEMICAL HAZARDS

STRONG OXIDIZING AGENT. CORROSIVE. Mix only with potable water at 60 - 80°F. Product must be diluted in accordance with label directions prior to use. The product is not combustible; however, at temperatures exceeding 156°F, decomposition occurs releasing oxygen. The oxygen release could initiate combustion. Never bring this product into contact with other sanitizers, cleaners or organic substances.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal. **PESTICIDE STORAGE:** Store only in original container. Keep this product under locked storage sufficient to make it inaccessible to children or persons unfamiliar with its proper use. **PESTICIDE DISPOSAL:** Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste Representative at the nearest EPA Regional Office for guidance. **CONTAINER HANDLING:** Non-Refillable Container. Do not reuse or refill this container. Triple rinse container promptly after emptying. Triple rinse as follows: Fill the container ¼ full with water and recap. Shake for 10 seconds. Drain for 10 seconds after the flow begins to drip. Follow Pesticide Disposal instructions for rinsate disposal. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration.